


Since many of the items lack a specific page number, the page number displayed online refers to the sequentially created number each item was given upon cataloging the materials.



In Eulhisel ruderalis Europ. ... Fr 1826 CECRCE ENGELMANN PAPERS MISSOUPI BOTANICAL GARDEN

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN

MISSOURI
BOTANICAL
GARDEN

Cuscuti astyla Caule funicalieni, glomembio zessilite racemose - spicatis, floribus singret. sesselibres sportitis. Tubo vorolle ogstridries calquis tegenenta ovata obliga parineta enetis vix excedente laviniis limbi orati tromavis incis structures to convavis incis triangularibus sussicilies, squamis tifisti ælæformibus orendata-fireline fancem atte sut more romple ste ng attingentitus; styli, coulities submillis, stigmete incrassation bilobo antheris brevore con lles ellipprometer circumsormant sapra capsulam proma calyptra marescente. John Mickets near Helenendorf, Georgia caucas. F. Hohenacken colle. 1. Um. itin. 1838. Currita omonogyna Un. itin collect. Holenal. Chorry monogra Juy 178. Very near G. monogyna, with which it has been impounded; but well distinguish by the smaller flowers, which are only half as large. The shorter telo af the corolle which handy exceeds the caly, The crembated lobes of the corolla. The ovate Frangulan not linear service of ignat, which reaches only to the lowest the stamina and Docs not come to the same level with their port obtuse capsule. _ The soules are as in the german form of C. monogyna bifit or wing like provesses on both with of the filaments, but they wears nearly up to the authors and are through of a firmer le Xtine. MISSOURI

BOTANICAL

GARDEN

copyright reserved

cm

Evica herbares L. , savat, lis Salisb German. auch Europauch Tarelad in Bohn MISSOURI BOTANICAL copyright reserved GARDEN

cm

Cuscula monogyna Vahl Caule funiculari glomemles sessitions varemoso-spication, floribus songue jub ics : Ci Gus Bester of elio - 5 pottilis tube corollae oglindrico, calycis segmente. ovata obtasa integris longe ex redente lavinias integris levetis, queterne longioni antheris obtas obtas inscrettes the Vier de longioni antheris oblongo lineambres subsensilis ous. squamis med im tubur, non supermitibles stylis coelits, ovarian vonice om aequantibles, stigmate siconasti tills authors non excedentibus; appell concila super capsulan magnam conicam cirrumsissum a lyitima, antheris I divisionen former subsens Colons squants Acuissimis beninger finhja Zej. Blaica i Agentina antheris in tuto corollae sub face statisticantilistic conveyent bus industris suffer stifitetie, squamis fermiosilus cratis orenulato-findriatis Enter Europe Silesia on the Oder on Salives etr Asia (Ming et); mr B. on the Altri (Levelus) my sperim on Annato! The species is remarkable in many other arrows to it has the thickets others of the lumpscan and or north american species tage finil, its stampers are inserted Jolow the face, in the tube itself. and the styles are connected into one and the Division can only be - observed in the slightly rapitate 2 lober stig zna Parepinent with of the capsule will opens transversely at lane, like the Currente with filiform obigines. The disept ment with remains with the bree of the regarder on the persister only is repeated from the upper the upper part of the appelle with a broad rim while the pursues the part of the disciplement of all Currentage with filiform stigmes is bilobed or obsordate leaving a part win the top part of 4

All The German spersmens examined to me have the menty the author, nearly zessile below the División of the Corolla! and have very this bifit scales on when the wally Vappendayes of the on both sides of the followend where is is menged into the substance of the corolle. are not connected in their free portion Do as to form one scale. __ . Affinished Var B. is ditinguishe by its brown to the souler of C. Guithy men which dre even sometimes grown together. and also by the longer filaments which inserted even lower in the take of the words though und, and are horizontely turned inwand to try the erect sign anthers closely around the -stigme. My sperime growing on Anndo is another proof of The possibility of Currette grand clours Munco. unon relations of E 7 7 1 30 · Desperon 21,40) 16/1/1/2


0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN

MISSOURI
BOTANICAL
GARDEN

exallato perifleris permetti, 3. Cusinta gamosty la Carle finiculari, orassissimme glomentis Del sessilibres, racemoso-spiratis, floribes singules breviter pedicellatis (s. sessilibres) 5 patitis, tabs corollar gylindren entydis lægententij ad lægin sepulis in dorso verricosis breviore, lacinis limbi Vosatas obtusas bis longiore.
antheris Tovatis Attaces, to bus, filamenti, latis tubo adratis, utmingue squamis alse formibus, distinctis, savius ef tulafut coalilis, dentates instructis, ovans globoso, stylis oraalitis, sligmations
glotapitator, trocis Capsula (magna) ovata corolla marrestente calquesta, circumstissa Along the guadloupe Citolo and other rivers near San Antonio Texas L'Lewhermen Aug & Septembe 1846 Colombing over thikely and smill frees also on the Colombia new La Grange (In Wright, - Leed yellow Entryor 1/4 coverner (and albamen) - albamen horny, rear Something like two scales - cotyledong! The roily of embrye not always regular.



Cuscuta elation Chois Cymndflor, Walkich not Hund Comb funioulari.

Thribus singulis pediallates . 5 partity tuby sorollae oplindstrom calquis segmentis ovato sub rotundad leviter cremelates, verneous - carinates el Carinis Cimbi onte obtessionsonlig petentity (Demun reflex is) questing quedrupla super toler longiore, antheris oblongs lineaniber subsemilier squamis finbriatis, medium tubus vix attingentle, ovario conice stylis bis connates incrassatis elongatis, apricen versag attenuati, Cynla Hes: Repairl, wallih. Hoteling for the West of the And of natural Science, Miles The flowers bear a semakett resemblance to those of C. chilensis, they have but The same shape a o the same or i little largen Tige. Et the onlyx is very Sifferent. This romes very men to that of Overlegerige i shape e sige it is also irendete or rather? alittle orliate on the many has and carrinate in the middle but the owner of the Sistinguish The word drows but by Ohning monge p. 177


wh is my small sperimen I want find the stem rough, It lobes of the corolle are with MISSOURI

BOTANICAL

GARDEN


copyright reserved

cm




Cusanta predicellata Del One repillaceo ramos, glomentes susilibres floribres singulais pedirellatio, out s'partitis talproor las campanulato segmenter calins Arianguli-ovata avet paulo supermete levini i limbi orates avules s. breviter abrupte patelas aequante; slamisibus Culo Isabi breviori bus filamente filiformates atteris orato orbination squamis truncatis supe bifides, aprice emargiants. s. Aremilatio; offis run strymatilen, Ailiformibus ovanis conico-brevionibus (2)
capsula commensorissa conolla marerente mundata Altri, Ledebour, my grevianen om Jalien. glomentes witheffer (5-5) flowers. flower freez small on periodle of the own Ength as of a very thin texture, It is menty retailed with Careline but has a much smaller flower shorter

0 1 2 3 4 5 6 7 8 9 10 MISSOURI BOTANICAL GARDEN



b. Con the stillish capsulam marres rente · Licht matiti d. Horolle squamis substamineis destitute. 4. C. desquamata 13. com. og. substan. pmedita * Colis corolle membetis -.5. C. arabica Fresen * K. Colis cor inlegnis † lobis corollar 4 partitar patentiles 6. C. minima ## Col. cor. 5 partitu subercotis

S. raule simplisi, corolla subirdusa 7. C. Cysilinum Was 5.8. combe namore conste exserte 8. C. Ennaviens reserved 33. cravio minore tabun corollac non replente stigmation mm stylis aquelibres ovario å. ordyre tala corollae sollaenione formione corolla cylindria Colis arminitis 9. C. Epithymum 2 6. only re tubun corollae longe superante 10. C. Coluteae a. calque profunde 5 partito 11. C. oupulatu De men breviter 5-lobo



0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

MISSOURI

BOTANICAL

GARDEN

Casatre stigmete capitate Il Meograe A Chlorocanneau 6. Corolla ad bisis, capsula marres rente 19 C. Volygonome & florite sul 4 - partitis. B floribus 5 partitis * lobis algeis orbiculato-obtasis 20. C. pentagona phinie. * lober objet acuter s'acuminate, Titglis ovanium subaquantibus. 9. calyce vermesos aspero 24 C verrue. 33 orlyre levi temi 22 C umbella tt. styles ovario melte longionets 23. Californ Bound carpeal ovaris stylopedi coronets
plus ministre con co capsul firmiore a tely in membrane ex marginile, integris Misso d. calyvibus monosepalis 4-5 portit, * lobis corollae oblusis † Capsula sulglobora. S. labis corollee tales brevioribes

Este ovatis

Corolle et brevioribes

pursistate 24 C. valgivaga \$\$ Corolle ad agricane 25. C. Comariensis &\$ Colis corolla tubum aquantity Comi - oblingis 26 C. Saururi ++ Capsula comica 5. nostrata 27. C. 0 x y carpa ** Color corollar cercitis t corolla tubulosa 28. C. chilensis Re ++ corolla exampement. 28. C coryenboxa. B. ratyribus 5 septes squamis sepelinis 2-7 printette datis. * squamis calyrinis adpressis † theyer tubun corolla subagaite 30 C. adpressa It ralge take rosolle Ingle Gresson 30. C. comparte ** squamis organis squamoni 32 C. glomenta 6. vorolla carnosa margine gorenulato L' Hontes 4 partités, oquanis minutes 33. C. Coryli B. Hen bus Spartite, squamis magnis. * calque d'après tubles, corolles aequante superante 34. C. Menropety * X- calyon his provide below of the 35 Chispidal MISSOURI

BOTANICAL copyright reserved cm GARDEN Cm Cm Cusmite stigmete capitate Il Meograe A Chlorocarpiere 6. Corolla at bisin oupsulue manrescente & lobis rulyonis orbinlatio obtains 19. C. pentagona B. Cobis cal. acutis s. accumination * Ttylis ovarium subacquent, En Tralyre lawi, Temus \$. floribus subsentibus, stylis 20. C. Polygonorum \$. fl. petioletis, stylis plijom. 21. C. umbellet. Ton tt algoe vermoor aspero 22. Premiera * * slylis ovenis much long with 23. C. culifornica Bound carpeal ovaris tylepedi coronets
plus mines ve con co, capsul, firmioge a tely in membrane ex margilite, integris Misso d. calyvibus monosepulis 4-5 partiti, * lobis covollae oblusis of Capsula sulglobosa, S. labis corollee tales brevious bus

Este ovatis

Corolle ad brein capsular 24 C. valgivaga

persistent 24 C. valgivaga tt Call at anicem

